

Lactobacillus plantarum FNCC 0137 fermented red Moringa oleifera exhibits protective effects in mice challenged with Salmonella typhi via TLR3/TLR4 inhibition and down-regulation of proinflammatory cy

by Mm Riyaniarti Estri Wuryandari, Et Al.

Submission date: 08-Dec-2022 01:27PM (UTC+0700)

Submission ID: 1975094123

File name: TLR_-_Maria_Wuryandari.pdf (2.77M)

Word count: 7115

Character count: 39264



Contents lists available at ScienceDirect

Journal of Ayurveda and Integrative Medicine

journal homepage: <http://elsevier.com/locate/jaim>



Original Research Article

Lactobacillus plantarum FNCC 0137 fermented red *Moringa oleifera* exhibits protective effects in mice challenged with *Salmonella typhi* via TLR3/TLR4 inhibition and down-regulation of proinflammatory cytokines



MM Riyaniarti Estri Wuryandari^{a,*}, Mochammad Fitri Atho'llah^b, Rizky Dzariyani Laili^c, Siti Fatmawati^d, Nashi Widodo^{b,f}, Edi Widjajanto^e, Muhaimin Rifa'i^{b,f,**}

^a Department of Biology, Faculty of Technology and Health Management, Institut Ilmu Kesehatan Bhakti Wiyata, 64114, Kediri, East Java, Indonesia

^b Department of Biology, Faculty of Mathematics and Natural Sciences, Brawijaya University, 65145, Malang, East Java, Indonesia

^c Department of Nutrition, Sekolah Tinggi Ilmu Kesehatan Hang Tuah Surabaya, 60244, Surabaya, East Java, Indonesia

^d Department of Food Sciences and Technology, Faculty of Agricultural Technology, Brawijaya University, 65145, Malang, East Java, Indonesia

^e Faculty of Medicine, Brawijaya University, 65145, Malang, East Java, Indonesia

^f Center of Biosystem Study, LPPM of Brawijaya University, 65145, Malang, East Java, Indonesia

ARTICLE INFO

Article history:

Received 10 August 2021

Received in revised form

2 October 2021

Accepted 22 October 2021

Available online 10 December 2021

Keywords:

Inflammation

Lactobacillus

Red moringa

Regulatory T cells

Salmonella

ABSTRACT

Background: *Salmonella typhi* is a foodborne pathogenic bacterium that threatens health. *S. typhi* infection exacerbated the antibiotic resistance problem that needs alternative strategies. *Moringa oleifera* possesses anti-inflammatory and antimicrobial effects. However, there is a lack of information about the pharmacological value of red *M. oleifera*. The fermentation of red *M. oleifera* leaves extract (RMOL) is expected to add to its nutritional value.

Objective: The present study aimed to evaluate non-fermented RMOL (NRMOL) and fermented RMOL (FRMOL) effects on *S. typhi* infection in mice.

Materials and methods: Female Balb/C mice were randomly divided into eight groups. The treatment groups were orally administered with NRMOL or FRMOL at doses 14, 42, and 84 mg/kg BW during the 28 days experimental period. Then *S. typhi* was introduced to mice through intraperitoneal injection except in the healthy groups. The NRMOL or FRMOL administration was continued for the next seven days. Cells that expressed CD11b⁺TLR3⁺, CD11b⁺TLR4⁺, CD11b⁺IL-6⁺, CD11b⁺IL-17⁺, CD11b⁺TNF- α ⁺, and CD4⁺CD25⁺CD62L⁺ were assessed by flow cytometry.

Results: Our result suggested that NRMOL and FRMOL extracts significantly reduced ($p < 0.05$) the expression of CD11b⁺TLR3⁺, CD11b⁺TLR4⁺, CD11b⁺IL-6⁺, CD11b⁺IL-17⁺, and CD11b⁺TNF- α ⁺ subsets. In contrast, NRMOL and FRMOL extracts significantly increased ($p < 0.05$) the expression of CD4⁺CD25⁺CD62L⁺ subsets. NRMOL at dose 14 and 42 mg/kg BW was more effective compared to FRMOL in reducing the expression of CD11b⁺TLR3⁺, CD11b⁺TLR4⁺, and CD11b⁺TNF- α ⁺ subsets.

Conclusion: Our findings demonstrated that NRMOL and FRMOL extracts could be promising agents for protection against *S. typhi* infection via modulation of TLR3/TLR4, regulatory T cells, and proinflammatory cytokines.

© 2021 The Authors. Published by Elsevier B.V. on behalf of Institute of Transdisciplinary Health Sciences and Technology and World Ayurveda Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Salmonella enterica serovar typhimurium (*Salmonella typhi*) is a Gram-negative bacterium that is an important foodborne pathogen with a worldwide distribution. *S. typhi* has received extensive attention due to its harmful effect on humans and animals,

* Corresponding author.

** Corresponding author.

E-mail addresses: mm.riyaniarti@iik.ac.id (M.R.E. Wuryandari), rifa123@ub.ac.id

(M. Rifa'i).

Peer review under responsibility of Transdisciplinary University, Bangalore.

<https://doi.org/10.1016/j.jaim.2021.10.003>

0975-9476/© 2021 The Authors. Published by Elsevier B.V. on behalf of Institute of Transdisciplinary Health Sciences and Technology and World Ayurveda Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

including abdominal pain, acute diarrhea, fever, nausea, vomiting, and sometimes lethal septicemia [1,2]. Interestingly, *S. typhi* infection in humans and mice produces many similar symptoms and hence mice are the most widely used animal model for studying *Salmonella* infection [3,4]. *Salmonella* infection was reported to affect around 11–21 million people worldwide, and approximately 128–161 thousand people die annually, with South-East Asia being one of the regions with highest case frequency [5].

As a first-line of defense, body initiates an acute inflammatory response in response to *S. typhi* infection. This response is executed via direct interaction of *S. typhi* with host cells such as macrophages or dendritic cells (DCs), followed by intensifying the inflammatory response, mainly in the liver, spleen, lungs, and intestines [6,7]. During bacterial infection, macrophages expressed Toll-like receptors 4 (TLR4), which primarily recognizes the lipopolysaccharide (LPS) as the major component of the cell wall of *S. typhi*. This recognition subsequently stimulates TLR4 signaling pathways, such as NF κ B, to produce proinflammatory cytokines, resulting in increased systemic cytokine production and septic shock in the later stages [8–10].

In the earliest phase of infection, both expressions of TLR3 and TLR4 in macrophages were increased to overproducing proinflammatory cytokines. These circumstances provide a survival advantage during bacterial elimination by producing reactive oxygen and nitrogen species, resulting in increased oxidative stress [11–14]. However, chronic inflammation and oxidative stress can promote tissue damage [15,16]. Therefore, the development of alternative options for the treatment or prevention that will reduce tissue damage caused by *S. typhi* and its virulence are urgently needed since *S. typhi*, and other bacteria become multi-resistant to antibiotics [17].

Moringa oleifera Lam. (Family: Moringaceae) originated from the Himalayas and was distributed almost worldwide in tropical and subtropical countries. *M. oleifera* is commonly known as horseradish tree, drumstick tree, or kelor (Indonesian). *M. oleifera* is also considered the magic tree or tree of life due to its abundant macro-, micro-, and phytonutrients [18,19]. Accumulating evidence showed that *M. oleifera* leaves extract (MOL) possessed antioxidant [20], anti-diarrheal [21], anti-inflammatory [22], and antimicrobial activities [23] due to its richness in flavonoids, flavanol glycosides, glucosinolate, isothiocyanate, phenolic acid, terpene, alkaloid, and sterol contents [24,25] However, the extract from MOL is not palatable [26].

It has been long recognized that utilization by microorganisms has been commonly used strategy to improve the functional properties of the plant [27,28]. The fermentation of MOL generate a sweet aroma that may increase its appeal and palatability [29]. Besides, MOL fermented by *Lactobacillus plantarum* alters its taste, pH, and viscosity [30]. Interestingly, fermentation of MOL with *L. plantarum* reduces its phytate and raffinose content and enhances its peptic digestibility and radical scavenging activity [26,31]. Further, milk fermented with *L. plantarum* demonstrated beneficial effects against *S. typhi* infection [32].

Although many studies have reported MOL's function and its product as an anti-inflammatory agent, there is still a lack of information about red MOL as anti-inflammation in the context of improving specific host immune function related to *S. typhi* infection. Red *M. oleifera* is not very popular as green *M. oleifera*. In contrast, the red *M. oleifera* is frequently used by local tribes in Southeast Sulawesi, Indonesia, as traditional medicine for curing various diseases than the green *M. oleifera* [33]. Herein, we provide evidence that red *M. oleifera* could restore naïve regulatory T cells and TLR3/TLR4 expression that affect proinflammatory cytokines, i.e., interleukin (IL)-6, IL-17, and TNF- α in macrophages of mice infected with *S. typhi*. Red Moringa may have anti-inflammatory activity through modulation of host-immune response in mice challenged with *S. typhi*.

2. Material and methods

2.1. Plant material and identification

The fresh portions of red *M. oleifera* leaves (leaves, seeds, flowers, and roots) were obtained from Sampang, Madura, East Java, Indonesia, during July 2017. The plant specimens then were authenticated and deposited at Purwodadi Botanic Garden, Indonesian Institute of Sciences, Pasuruan Indonesia, with the voucher specimen numbers 1051/IPH.06/HM/VIII/2017.

2.2. *Moringa oleifera* preparation

Red *M. oleifera* leaves were washed three times using distilled water. Then the leaves were air-dried for 72 h, followed by drying in an oven at 40 °C for 3 h. Dried leaves of red *M. oleifera* were ground to obtain a powder. The leaves powder (200 g) then macerated with 2 L 70% ethanol for three consecutive days, continuously shaking at 125 rpm for 1 hour per day. The red *M. oleifera* leaves extract (RMOL) was filtered using Whatman No. 1 paper and then concentrated using a rotary evaporator (IKA RV10).

2.3. *Moringa oleifera* fermentation

L. plantarum FNCC 0137 was received from the Food and Nutrition Study Center, Gadjah Mada University. *L. plantarum* FNCC 0137 was prepared using the MRS broth medium and incubated at 37 °C for 72 hours followed by centrifugation for 20 min at 4 °C. MOL extract at room temperature was fermented with 1×10^8 CFU/g *L. plantarum* FNCC 0137, followed by incubation at 37 °C for 120 h [31]. The fermentation product of RMOL (FRMOL) was supplemented with 10% sucrose and 0.5% NaCl prior to freeze-drying [34].

2.4. Experimental animal

Female *Balb/C* mice were obtained from the Institute of Biosciences, Brawijaya University. Female *Balb/C* mice with 25–30 g weight and six weeks old were housed at Animal Facility, Faculty of Agricultural, Brawijaya University. Mice were allowed to consume food and water *ad libitum*. Mice were acclimatized for seven days at 25–26 °C, RH 60%, and with a 12 h light/dark cycle.

2.5. Non-fermented and fermented MOL administration

After one-week acclimatization, forty female *Balb/C* mice were randomly divided into eight groups (n = 5):

- Group I = Healthy mice without *S. typhi* injection (HM)
- Group II = *S. typhi* only without additional administration (*S. typhi*)
- Group III = NRMOL 14 mg/kg BW + *S. typhi* (NRMOL – 14)
- Group IV = NRMOL 42 mg/kg BW + *S. typhi* (NRMOL – 42)
- Group V = NRMOL 84 mg/kg BW + *S. typhi* (NRMOL – 84)
- Group VI = FRMOL 14 mg/kg BW + *S. typhi* (FRMOL – 14)
- Group VII = FRMOL 42 mg/kg BW + *S. typhi* (FRMOL – 42)
- Group VIII = FRMOL 84 mg/kg BW + *S. typhi* (FRMOL – 84)

Both NRMOL and FRMOL were given orally for 28 consecutive days. On the 29th day, mice except for healthy mice groups were intraperitoneally infected with 1×10^7 CFU/ml *S. typhi* for the next seven days.

2.6. *S. typhi* confirmation

After 24 h, *S. typhi* infection was confirmed through collected blood via vena caudalis. Briefly, 50 μ L blood was collected and

450 μ L of 0.9% sterile NaCl was added to it. The mixture was then planted in Luria broth medium and incubated for 24 h at 37 °C with 120 rpm. Followed by isolation from Luria Broth and inoculation in Salmonella selective medium, Xylose-Lysine-Deoxycholate (XLD). The colony of Salmonella is characterized by a colony that formed dark nuclei with a transparent colony. Further confirmation for Salmonella was carried out using the catalase test. Briefly, one ose from cultured Salmonella was dipped in object-glass containing H₂O₂. formation of bubble was confirmed presence of Salmonella [35,36].

2.7. Splenocyte isolation

Mice were abstained from food overnight with free access to water before being sacrificed through cervical dislocation. Spleen was collected, washed, and crushed to obtain a single-cell suspension. The homogenate was then centrifuged at 2500 rpm for 5 min at 10 °C. The supernatant was discarded, and the pellet obtained was added with 1 mL PBS. Homogenates was aliquoted into 1.5 mL tubes and centrifuged to remove supernatant. The pellet was then stained with appropriate antibodies for flow cytometry analysis.

2.8. Flow cytometry staining

The antibodies used for extra- and intracellular staining were purchased from BioLegend (San Diego, CA, USA) and prepared following routine laboratory procedures [37]. Regulatory T cells (Tregs) were marked by the combination of fluorescein isothiocyanate (FITC) anti-mouse CD4 (clone GK1.5), phycoerythrin (PE) anti-mouse CD25 (clone 3C7), and PE-Cy5 anti-mouse CD62L (clone MEL-14) as cell surface antibodies for 30 min at 4 °C in the dark. Macrophages were identified by cell-surface antibodies FITC anti-mouse/human CD11b (clone M1/70) for 30 min at 4 °C in the dark, followed by a wash step and added cytofix/cytoperm for 20 min at 4 °C. The supernatant was discarded after centrifugation, then the pellet obtained was stained with PE anti-mouse TLR3 (clone 11F8), PE-Cy7 anti-mouse TLR4 (clone SA15-21), PE anti-mouse IL-6 (clone MP5-20F3), PE-Cy7 anti-mouse IL-17A (clone TC11-18H10.1), and PE anti-mouse TNF- α (clone MP6-XT22). Cells were resuspended in PBS, and a total of 10,000 cell events were acquired using FACS Calibur™ at a low or medium rate. The cells' population was then according to the stained used for further analysis. Cell subsets analysis was performed using FlowJo v10 for Windows (FlowJo LLC, Ashland, OR), following the previously validated protocol [38].

2.9. Statistical analysis

One-way analysis of variance (ANOVA) was used to determine statistically significant differences between groups (p -value < 0.05). Tukey HSD test was used for multiple comparisons and post hoc analysis. Analyses were performed using GraphPad Prism 8 (GraphPad Software Inc, La Jolla, CA).

3. Result

3.1. NRMOL restores CD11b⁺TLR3⁺ and CD11b⁺TLR4⁺ subsets better than FRMOL in mice challenged with *S. typhi*

In the present study, we try to assess the NRMOL and FRMOL efficacy on the expression of CD11b⁺TLR3⁺ and CD11b⁺TLR4⁺ subsets (Fig. 1A–D). Our result suggest that NRMOL is more effective than FRMOL to reduce CD11b⁺TLR3⁺ and CD11b⁺TLR4⁺ subsets after *S. typhi* challenge. Based on dot plot analysis, CD11b⁺TLR3⁺

subsets (Fig. 1A) and CD11b⁺TLR4⁺ subsets (Fig. 1C) were increased in mice challenged with *S. typhi*. Both NRMOL and FRMOL administration significantly reduced ($p < 0.05$) the CD11b⁺TLR3⁺ subsets (Fig. 1B) and CD11b⁺TLR4⁺ subsets (Fig. 1D) compared to mice challenged with *S. typhi* only. Interestingly, NRMOL at doses 14 and 42 mg/kg BW more effective than FRMOL to reduce CD11b⁺TLR3⁺ and CD11b⁺TLR4⁺ subsets in mice challenged with *S. typhi*.

3.2. NRMOL and FRMOL declines proinflammatory cytokines in mice challenged with *S. typhi*

Proinflammatory cytokines are the main end-product of inflammatory processes. The present study data suggest that NRMOL is more effective than FRMOL to reduce CD11b⁺IL-6⁺ (Fig. 2A), shared the same better effect to reduce CD11b⁺IL-17⁺ subsets (Fig. 2C), and NRMOL reduces CD11b⁺TNF- α ⁺ subsets more effective FRMOL after *S. typhi* challenge (Fig. 2E). Based on dot plot analysis, CD11b⁺IL-6⁺, CD11b⁺IL-17⁺, and CD11b⁺TNF- α ⁺ subsets were increased in mice challenged with *S. typhi*. Both NRMOL and FRMOL administration significantly reduced ($p < 0.05$) the CD11b⁺IL-6⁺, CD11b⁺IL-17⁺, and CD11b⁺TNF- α ⁺ subsets compared to mice challenged with *S. typhi* only. Interestingly, NRMOL at doses 14 mg/kg BW is more effective than FRMOL to reduce CD11b⁺IL-6⁺ subsets in mice challenged with *S. typhi*. Surprisingly, NRMOL at doses 42 and 84 mg/kg BW reduced CD11b⁺IL-6⁺ subsets lower than healthy mice groups (Fig. 2B). Interestingly, NRMOL and FRMOL at doses 14 and 42 mg/kg BW reduce CD11b⁺IL-17⁺ subsets in mice challenged with *S. typhi* towards near healthy mice groups (Fig. 2D). NRMOL at doses 14 and 42 mg/kg BW is more effective than FRMOL to reduce CD11b⁺TNF- α ⁺ subsets in mice challenged with *S. typhi* (Fig. 2F).

3.3. NRMOL and FRMOL restores naïve regulatory T cells in mice challenged with *S. typhi*

NRMOL and FRMOL had similar efficacy to improve CD4⁺CD25⁺CD62L⁺ subsets after *S. typhi* challenge (Fig. 3A–B). Based on dot plot analysis, CD4⁺CD25⁺CD62L⁺ subsets were decreased in mice challenged with *S. typhi* (Fig. 3A). Both NRMOL and FRMOL administration significantly increased the CD4⁺CD25⁺CD62L⁺ subsets ($p < 0.05$) compared to mice challenged with *S. typhi* only (Fig. 3B). Interestingly, NRMOL and FRMOL restored CD4⁺CD25⁺CD62L⁺ subsets in mice challenged with *S. typhi* similar to healthy mice groups, except NRMOL dose 14 mg/kg BW.

4. Discussion

Salmonella constitutes a considerable health burden by causing acute gastroenteritis, contributing almost half of morbidity and mortality attributable to typhoid fever in low- and middle-income countries. Antibiotics are commonly used to treat Salmonellosis in humans, however because of antibiotic resistance antibiotics are not entirely effective in combatting *Salmonella* [39]. Developing new products from natural plants can be an effective strategy however it is still challenging for many researchers for past few decades. Fermentation is a useful method for improving food products' biological properties, promoting the beneficial effects on health. Numerous evidences reported that fermentation using probiotics can be an alternative option to combat *Salmonella* [32,40].

Chronic *Salmonella* infection occurs when the host fails to completely clear bacteria from the body. An impaired immune response or gut microbiota disruption may be associated with the host's inability to clear *Salmonella* [41]. Our result demonstrated




Fig. 1. Effect of NRMOL and FRMOL on CD11b⁺TLR3⁺ and CD11b⁺TLR4⁺ subsets in mice challenged with *S. typhi*. (A) The dot plot analysis of CD11b⁺TLR3⁺ subsets in spleen. (B) NRMOL reduced CD11b⁺TLR3⁺ subsets more effective FRMOL on mice challenged with *S. typhi*. (C) The dot plot analysis of CD11b⁺TLR4⁺ subsets in spleen. (D) NRMOL reduced CD11b⁺TLR4⁺ subsets more effective FRMOL on mice challenged with *S. typhi*. The different letter considered significantly different between each group ($p < 0.05$) by post hoc test using Tukey's HSD test. HM = Healthy Mice; NRMOL = Non-fermented red *M. oleifera* Leaves Extract; FRMOL = Fermented red *M. oleifera* Leaves Extract; NRMOL-14 = *S. typhi* + NRMOL 14 mg/kg BW; NRMOL-42 = *S. typhi* + NRMOL 42 mg/kg BW; NRMOL-84 = *S. typhi* + NRMOL 84 mg/kg BW; FRMOL-14 = *S. typhi* + FRMOL 14 mg/kg BW; FRMOL-42 = *S. typhi* + FRMOL 42 mg/kg BW; and FRMOL-84 = *S. typhi* + FRMOL 84 mg/kg BW.

that *S. typhi* elicits an immune response through an increase in expression of CD11b⁺ TLR3⁺ and CD11b⁺TLR4⁺ subsets. At the earliest infection phase by *Salmonella*, macrophages secrete proinflammatory cytokines, such as interferon- β (IFN- β), through TLR3 and TLR4 signaling pathways [13]. TLR3 and TLR4 share the same adaptor molecule, toll/interleukin-1 receptor (TIR) domain-containing adapter-inducing IFN- β (TRIF), which activates NF κ B, interferon regulatory factor 3 (IRF3), and MAP kinase leading to the type I IFN transcription [42]. IFN will protect the host through antimicrobial autophagy by macrophages, while, the over-expression of IFN may result in impaired bacterial clearance [43]. *Salmonella* infection would worsen if the host cannot remove *Salmonella* from the body completely because the *Salmonella* present in the body will cause chronic inflammation and will also produce toxins that will induce mucosal damage [44]. A previous study reported that presence of *S. typhi* was observed in the caecum at 7-days post-infection. Moreover, the serum concentration of proinflammatory cytokines, including TNF- α and IL-6, was higher compared to control, indicating that *S. typhi* generated an inflammatory response during infection and led to caecum damage [45].

In the present study, NRMOL and FRMOL inhibited TLR3/TLR4 compared to the increased expression of TLR3/TLR4 detected in the *S. typhi* group. The bioactive compounds from MOL have improved host immunity and inhibited LPS induced inflammatory responses. LPS is well-known as the primary ligand for TLR4

signaling activation [46,47]. Some evidence suggests that FRMOL could be a potent immunomodulator [35,36,48]. We would like to propose two possible mechanisms through which MOL could downregulate the TLR3 expression. First, the bioactive compounds of MOL influence TLR3/TRIF-dependent pathways. At various degrees, flavonoids can alter TLR pathways. Flavonoids can impact both TLR gene and cell membrane expression, which are both directly related to TLR functionality [49]. Luteolin and quercetin, present in MOL, were demonstrated to suppress the TRIF-dependent pathway. Interestingly, both luteolin and quercetin have a C₂-C₃ double bond in the carbonyl-containing C-ring, which is important for TANK-binding kinase 1 (TBK1) inhibition [50]. TRIF activates TBK1, phosphorylates IRF3 and induces the IFNs expression [13,42]. Further, luteolin's chemical structure, which has two hydroxyl groups at position 3' and 4' in B-ring, was considered an important factor to target TLR3 signaling pathways [50]. Second, the bioactive compounds of MOL may contribute indirectly through tissue repair. Our previous study revealed that FRMOL has a hepatoprotective effect via Nrf2 signaling pathways in mice challenged with *S. typhi*. FRMOL administration also repairs hepatocyte damage through reduced necrotic cells [51]. TLR3 could be activated through extracellular dsRNA released from damaged tissue [52]. The diminished expression of TLR3 by NRMOL or FRMOL may be caused by repairing tissue damage, which reduces the TLR3 main ligand.




Fig. 2. Effect of non-fermented MOLE and fermented MOLE on proinflammatory generation in mice challenged with *S. typhi*. (A) The dot plot analysis of CD11b⁺IL-6⁺ subsets in spleen. (B) NRMOL reduced CD11b⁺IL-6⁺ subsets more effective FRMOL on mice challenged with *S. typhi*. (C) The dot plot analysis of CD11b⁺IL-17⁺ subsets in spleen. (D) NRMOL and FRMOL reduced CD11b⁺IL-17⁺ subsets on mice challenged with *S. typhi*. (E) Dot plot analysis of CD11b⁺TNF-α⁺ subsets in spleen. (F) NRMOL reduced CD11b⁺TNF-α⁺ subsets more effective FRMOL on mice challenged with *S. typhi*. The different letter considered significantly different between each group ($p < 0.05$) by post hoc test using Tukey's HSD test. HM = Healthy Mice; NRMOL = Non-fermented red *M. oleifera* Leaves Extract; FRMOL = Fermented red *M. oleifera* Leaves Extract; NRMOL-14 = *S. typhi* + NRMOL 14 mg/kg BW; NRMOL-42 = *S. typhi* + NRMOL 42 mg/kg BW; NRMOL-84 = *S. typhi* + NRMOL 84 mg/kg BW; FRMOL-14 = *S. typhi* + FRMOL 14 mg/kg BW; FRMOL-42 = *S. typhi* + FRMOL 42 mg/kg BW; and FRMOL-84 = *S. typhi* + FRMOL 84 mg/kg BW.

The data from the present study demonstrated that NRMOL inhibits proinflammatory cytokines, IL-6, IL-17, and TNF-α secreted by macrophages. These findings are in agreement with the previously published results by our group that inhibition of TLR3 and TLR4 as upstream of signaling pathways would give a beneficial implication. *S. typhi* infection amplifies an inflammatory response

and induces macrophages to secrete a huge number of proinflammatory cytokines [9]. TNF-α is essential for generating a systemic inflammatory response that leads to lethal shock [53]. Interestingly, TNF-α collaboration with IL-17 triggers other proinflammatory cytokines production [54]. A recent study reported that red *M. oleifera* contains higher quercetin than green *M. oleifera* [55].




Fig. 3. Effect of NRMOL and FRMOL on CD4⁺CD25⁺CD62L⁺ subsets in mice challenged with *S. typhi*. (A) The dot plot analysis of CD4⁺CD25⁺CD62L⁺ subsets in spleen. (B) NRMOL and FRMOL increased CD4⁺CD25⁺CD62L⁺ subsets on mice challenged with *S. typhi*. The different letter considered significantly different between each group ($p < 0.05$) by post hoc test using Tukey's HSD test. HM = Healthy Mice; NRMOL = Non-fermented red *M. oleifera* Leaves Extract; FRMOL = Fermented red *M. oleifera* Leaves Extract; NRMOL-14 = *S. typhi* + NRMOL 14 mg/kg BW; NRMOL-42 = *S. typhi* + NRMOL 42 mg/kg BW; NRMOL-84 = *S. typhi* + NRMOL 84 mg/kg BW; FRMOL-14 = *S. typhi* + FRMOL 14 mg/kg BW; FRMOL-42 = *S. typhi* + FRMOL 42 mg/kg BW; and FRMOL-84 = *S. typhi* + FRMOL 84 mg/kg BW.

Quercetin is reported to suppress the lethal shock caused by *S. typhi* infection. The hydroxyl groups of quercetin are important factors for delaying *S. typhi* action [53]. Our previous study also showed that FRMOL increased its total flavonoid content [35]. Our findings are also supported by a study that reported milk fermented with *Lactobacillus* reduced IL-17, TNF- α , and IL-6 in mice challenged with *Salmonella* [32,56]. Acurcio et al., 2017 proposed that the possible mechanism of a protective effect from *L. plantarum* is due to modulation of the host immune system [32]. Further, probiotic use could stimulate the anti-inflammatory cytokine expression, followed by reducing proinflammatory cytokines [57]. The anti-inflammatory properties of MOL have been reported to decline

NF κ B and proinflammatory cytokines expression in several animal models of inflammation [58,59]. Based on our findings, we assumed that both NRMOL and FRMOL might be useful to also treat other inflammatory diseases through the downregulation of inflammatory cytokines in the TLRs signaling pathway.

Our result also demonstrated that NRMOL and FRMOL restored the naïve Tregs. Tregs are essential to maintain peripheral immune tolerance to self-antigen. Tregs play an important role in controlling immune response during infection [60]. Tregs suppress the effector T cells by various mechanisms, mainly producing IL-10. The balance between Tregs and effector T cells reflects the immune homeostasis [61]. A previous study reported that quercetin and other flavonoids

could modulate the aryl hydrocarbon receptor (Ahr), which further influenced Tregs' activation via Foxp3, the main transcription factor for Tregs [62–64]. On the other hand, the activation of heme oxygenase-1 (HO1) via the Nrf2/HO-1 signaling pathway is reported to increase Foxp3 expression. Meanwhile, a previous study reported that FRMOL increased HO-1 expression via the Nrf2/HO-1 signaling pathway. Naïve Tregs (CD4⁺CD25⁺CD62L⁺ subsets) could be influenced by nutrients and improve their suppressive function and number [65–67]. Based on our result, we suggest that the bioactive compound in NRMOL and FRMOL may benefit from restoring naïve Tregs, which may implicate Th cell responses during *S. typhi* infection.

5. Conclusion

In summary, our finding showed that NRMOL efficiently reduced CD11b⁺TLR3⁺, CD11b⁺TLR4⁺, CD11b⁺IL-6⁺ and CD11b⁺TNF- α ⁺ compared to FRMOL. In addition, NRMOL and FRMOL displayed similar effect in reducing CD11b⁺IL-17⁺ and restoring naïve Tregs. The main goal of anti-inflammatory therapy is inhibition of TLR3/TLR4 and the proinflammatory cytokines followed by the recovery of naïve Tregs. Our study provides evidence that the anti-inflammatory properties of RMOL could become a promising supplement to treat *S. typhi* infection or might be other inflammatory diseases. Future studies are required for exploring, the molecular mechanisms of NRMOL and FRMOL.

Ethical approval

All experimental procedures were approved by the Animal Care and Use Committee of Brawijaya University with approval number: 829-KEP-UB. All the experiments were conducted following the Guide to the Care and Use of Laboratory Animals (National Institutes of Health, United States).

Source of funding

The grant was funded by two Ministry: Ministry of Finance and the Ministry of Research Technology Higher Education through the program namely Endowment Fund for Education (LPDP-BUDI DN).

Conflict of interest

None.

Author contributions

MM Riyaniarti Estri Wuryandari: Conceptualization, Funding acquisition, Methodology, Writing - Original draft preparation. **Mochammad Fitri Athoillah:** Formal analysis, Writing - Original draft preparation, Visualization. **Rizky Dzariyani Laili:** Data curation, Investigation. **Siti Fatmawati:** Data curation, Investigation. **Nashi Widodo:** Supervision, Writing - Review & Editing. **Edi Widjajanto:** Supervision, Writing- Reviewing and Editing. **Muhaimin Rifai:** Conceptualization, Resources, Supervision, Writing Review & Editing.

Acknowledgements

We thanks to Laboratory of Animal Physiology, Structure, and Development, and Laboratory of Microbiology, Brawijaya University for provide the facilities.

References

- Andritschke D, Dilling S, Emmenlauer M, Welz T, Schmach F, Misselwitz B, et al. A genome-wide siRNA screen implicates spire1/2 in SipA-driven *Salmonella typhimurium* host cell invasion. *PLoS One* 2016;11:e0161965. <https://doi.org/10.1371/journal.pone.0161965>.
- Coburn B, Li Y, Owen D, Vallance BA, Finlay BB. *Salmonella enterica* serovar typhimurium pathogenicity island 2 is necessary for complete virulence in a mouse model of infectious enterocolitis. *Infect Immun* 2005;73:3219–27. <https://doi.org/10.1128/IAI.73.6.3219-3227.2005>.
- Mian MF, Pek EA, Chenoweth MJ, Coombes BK, Ashkar AA. Humanized mice for *Salmonella typhi* infection: new tools for an old problem. *Virulence* 2011;2:248–52. <https://doi.org/10.4161/viru.2.3.16133>.
- Nilsson OR, Kari L, Steele-Mortimer O. Foodborne infection of mice with *Salmonella Typhimurium*. *PLoS One* 2019;14:e0215190. <https://doi.org/10.1371/journal.pone.0215190>.
- World Health Organization. Typhoid vaccine: WHO position paper - march 2018. *Wkly Epidemiol Rec* 2018;13:153–72.
- Dar MA, Ahmed R, Urwat U, Ahmad SM, Dar PA, Kushoo ZA, et al. Expression kinetics of natural resistance associated macrophage protein (NRAMP) genes in *Salmonella Typhimurium*-infected chicken. *BMC Vet Res* 2018;14:180. <https://doi.org/10.1186/s12917-018-1510-4>.
- Srikanth CV, Cherayil BJ. Intestinal innate immunity and the pathogenesis of *Salmonella enteritis*. *Immunol Res* 2007;37:61–77. <https://doi.org/10.1007/BF02686090>.
- Mogensen TH. Pathogen recognition and inflammatory signaling in innate immune defenses. *Clin Microbiol Rev* 2009;22:240–73. <https://doi.org/10.1128/CMR.00046-08>.
- Mathur R, Oh H, Zhang D, Park S-G, Seo J, Koblansky A, et al. A mouse model of *Salmonella typhi* infection. *Cell* 2012;151:590–602. <https://doi.org/10.1016/j.cell.2012.08.042>.
- Lestari SR, Atho'illah MF, Yi Christina, Rifa'i M. Single garlic oil modulates T cells activation and proinflammatory cytokine in mice with high fat diet. *J Ayurveda Integr Med* 2020;11:414–20. <https://doi.org/10.1016/j.jaim.2020.06.009>.
- Bihl F, Salez L, Beaubier M, Torres D, Larivière L, Laroche L, et al. Overexpression of toll-like receptor 4 amplifies the host response to lipopolysaccharide and provides a survival advantage in transgenic mice. *J Immunol* 2003;170:6141–50. <https://doi.org/10.4049/jimmunol.170.12.6141>.
- Deng S, Yu K, Zhang B, Yao Y, Wang Z, Zhang J, et al. Toll-like receptor 4 promotes NO synthesis by upregulating GCHI expression under oxidative stress conditions in sheep monocytes/macrophages. *Oxid Med Cell Longev* 2015;2015:1–11. <https://doi.org/10.1155/2015/359315>.
- Owen KA, Anderson CJ, Casanova JE. *Salmonella* suppresses the TRIF-dependent type I interferon response in macrophages. *mBio* 2016;7:e02051-15. <https://doi.org/10.1128/mBio.02051-15>. /mbio/7/1/e02051-15.atom.
- Zou J, Shankar N. Roles of TLR/MyD88/MAPK/NF- κ B signaling pathways in the regulation of phagocytosis and proinflammatory cytokine expression in response to *E. faecalis* infection. *PLoS One* 2015;10:e0136947. <https://doi.org/10.1371/journal.pone.0136947>.
- Arifah SN, Atho'illah MF, Lukiati B, Lestari SR. Herbal medicine from single clove garlic oil extract ameliorates hepatic steatosis and oxidative status in high fat diet mice. *Malays J Med Sci MJMS* 2020;27:46–56. <https://doi.org/10.21315/mjms2020.27.1.5>.
- Yu JH. Oxidative stress and inflammatory signaling in cerulein pancreatitis. *World J Gastroenterol* 2014;20:17324. <https://doi.org/10.3748/wjg.v20.i46.17324>.
- Whitchard JM, Gay K, Stevenson JE, Joyce KJ, Cooper KL, Omondi M, et al. Human *Salmonella* and concurrent decreased susceptibility to quinolones and extended-spectrum cephalosporins. *Emerg Infect Dis* 2007;13:1681–8. <https://doi.org/10.3201/eid1311.061438>.
- Gopalakrishnan L, Doriya K, Kumar DS. *Moringa oleifera*: a review on nutritive importance and its medicinal application. *Food Sci Hum Wellness* 2016;5:49–56. <https://doi.org/10.1016/j.fshw.2016.04.001>.
- Oyeyinka AT, Oyeyinka SA. *Moringa oleifera* as a food fortificant: recent trends and prospects. *J Saudi Soc Agric Sci* 2018;17:127–36. <https://doi.org/10.1016/j.jssas.2016.02.002>.
- Saini RK, Shetty NP, Prakash M, Giridhar P. Effect of dehydration methods on retention of carotenoids, tocopherols, ascorbic acid and antioxidant activity in *Moringa oleifera* leaves and preparation of a RTE product. *J Food Sci Technol* 2014;51:2176–82. <https://doi.org/10.1007/s13197-014-1264-3>.
- Misra A, Srivastava S, Srivastava M. Evaluation of anti diarrheal potential of *Moringa oleifera* (Lam.) leaves. *J Pharmacogn Phytochem* 2014;2:43–6.
- Xu Y-B, Chen G-L, Guo M-Q. Antioxidant and anti-inflammatory activities of the crude extracts of *Moringa oleifera* from Kenya and their correlations with flavonoids. *Antioxidants* 2019;8:296. <https://doi.org/10.3390/antiox8080296>.
- Bancesi A, Pinto MMF, Duarte E, Catarino L, Nazareth T. The antimicrobial properties of *Moringa oleifera* Lam. for water treatment: a systematic review. *SN Appl Sci* 2020;2:323. <https://doi.org/10.1007/s42452-020-2142-4>.
- Rani NZA, Husain K, Kumolosasi E. *Moringa* genus: a review of phytochemistry and pharmacology. *Front Pharmacol* 2018;9:108. <https://doi.org/10.3389/fphar.2018.00108>.

- [25] Leone A, Spada A, Battezzati A, Schiraldi A, Aristil J, Bertoli S. *Moringa oleifera* seeds and oil: characteristics and uses for human health. *Int J Mol Sci* 2016;17: 2141. <https://doi.org/10.3390/ijms17122141>.
- [26] Vanajakshi V, Vijayendra SVN, Varadaraj MC, Venkateswaran C, Agrawal R. Optimization of a probiotic beverage based on Moringa leaves and beetroot. *LWT - Food Sci Technol (Lebensmittel-Wissenschaft -Technol)* 2015;63: 1268–73. <https://doi.org/10.1016/j.lwt.2015.04.023>.
- [27] Atho'llah MF, Safitri YD, Nur'aini FD, Savitri RU, Rahayu S, Widyarti S, et al. Evaluation of glyceollin accumulation and antioxidant properties on soybean (*Glycine max* L.) through combination of different biotic elicitor and light. *Sci Study Res Chem Chem Eng Biotechnol Food Ind* 2019;20:199–208.
- [28] Feitosa PRB, Santos TRJ, Gualberto NC, Narain N, de Aquino Santana LCL. Solid-state fermentation with *Aspergillus niger* for the bio-enrichment of bioactive compounds in *Moringa oleifera* (moringa) leaves. *Biocatal Agric Biotechnol* 2020;27:101709. <https://doi.org/10.1016/j.bcab.2020.101709>.
- [29] Wang J, Cao F, Zhu Z, Zhang X, Sheng Q, Qin W. Improvement of quality and digestibility of moringa oleifera leaves feed via solid-state fermentation by *Aspergillus Niger*. *Int J Chem React Eng* 2018;16. <https://doi.org/10.1515/ijcre-2018-0094>.
- [30] Isnaeni I, Maulidina A, Kusumawati I, Setyawatie EM. Inhibitory activity of *Lactobacillus plantarum* ATCC 8014 fermented milk combined with aqueous extract of *Moringa oleifera* leaves against *Streptococcus mutans*. *J Res Pharm* 2019;23:701–10. <https://doi.org/10.12991/jrp.2019.179>.
- [31] Thierry NN, Léopold TN, Didier M, Moses FMC. Effect of pure culture fermentation on biochemical composition of *Moringa oleifera* Lam leaves powders. *Food Nutr Sci* 2013;4:851–9. <https://doi.org/10.4236/fns.2013.48111>.
- [32] Acurcio LB, Bastos RW, Sandes SH de C, Guimaraes AC de C, Alves CG, Reis DC dos, et al. Protective effects of milk fermented by *Lactobacillus plantarum* B7 from Brazilian artisanal cheese on a *Salmonella enterica* serovar Typhimurium infection in BALB/c mice. *J Funct Foods* 2017;33:436–45. <https://doi.org/10.1016/j.jff.2017.04.010>.
- [33] Sofyani WOW. Sistem klasifikasi kelor dalam etnobotani masyarakat wolio. *J Sosiol Walisongo* 2019;3:49–64. <https://doi.org/10.21580/jsw.2019.3.1.3488>.
- [34] Fonseca F, Cenard S, Passot S. Freeze-drying of lactic acid bacteria. In: *Wolkers WF, Oldenhof H, editors. Cryopreserv. Free-Dry. Protoc.*, vol. 1257. New York, NY: Springer New York; 2015. p. 477–88. https://doi.org/10.1007/978-1-4939-2193-5_24.
- [35] Laili RD, Martati E, Rifa'i M. Immunomodulator effect of *Moringa oleifera* leaves fermented by *Lactobacillus plantarum* FNCC 0137 on *Salmonella typhi* infected Balb/c mice. *Res J Pharm Technol* 2019;12:3595. <https://doi.org/10.5958/0974-360X.2019.00613.9>.
- [36] Fatmawati S, Laili RD, Wuryandari MRE, Martati E, Widyansih TD, Rifa'i M. Fermented ethanolic extract of *Moringa oleifera* leaves with *Lactobacillus plantarum* FNCC 0137 as immunomodulators on *Salmonella typhi* infected mice. *Res J Pharm Technol* 2020;13(12):1–6.
- [37] Safitri YD, Atho'llah MF, Nur'aini FD, Widyarti S, Rifa'i M. The effects of elicited soybean (*Glycine max*) extract on hematopoietic cells of high fat-fructose diet Balb/C mice model. *Jordan J Biol Sci* 2018;11:241–6.
- [38] Atho'llah MF, Safitri YD, Nur'aini FD, Widyarti S, Tsuboi H, Rifa'i M. Elicited soybean extract attenuates proinflammatory cytokines expression by modulating TLR3/TLR4 activation in high-fat, high-fructose diet mice. *J Ayurveda Integr Med* 2021;12:43–51. <https://doi.org/10.1016/j.jaim.2021.01.003>.
- [39] Gupta PD, Birdi TJ. Development of botanicals to combat antibiotic resistance. *J Ayurveda Integr Med* 2017;8:266–75. <https://doi.org/10.1016/j.jaim.2017.05.004>.
- [40] Wagner RD, Johnson SJ. Probiotic bacteria prevent *Salmonella* - induced suppression of lymphoproliferation in mice by an immunomodulatory mechanism. *BMC Microbiol* 2017;17:77. <https://doi.org/10.1186/s12866-017-0990-x>.
- [41] Zha L, Garrett S, Sun J. *Salmonella* infection in chronic inflammation and gastrointestinal cancer. *Diseases* 2019;7:28. <https://doi.org/10.3390/diseases7010028>.
- [42] Ruiz J, Kanagavelu S, Flores C, Romero L, Riveron R, Shih DQ, et al. *Front Cell Infect Microbiol* 2016;5. <https://doi.org/10.3389/fcimb.2015.00105>.
- [43] Manca C, Tsenova L, Bergtold A, Freeman S, Tovey M, Musser JM, et al. Virulence of a *Mycobacterium tuberculosis* clinical isolate in mice is determined by failure to induce Th1 type immunity and is associated with induction of IFN-. *Proc Natl Acad Sci Unit States Am* 2001;98:5752–7. <https://doi.org/10.1073/pnas.091096998>.
- [44] Song J, Gao X, Galán JE. Structure and function of the *Salmonella typhi* chimaeric A2B5 typhoid toxin. *Nature* 2013;499:350–4. <https://doi.org/10.1038/nature12377>.
- [45] Wu S-C, Chu X-L, Su J-Q, Cui Z-Q, Zhang L-Y, Yu Z-J, et al. Baicalin protects mice against *Salmonella typhimurium* infection via the modulation of both bacterial virulence and host response. *Phytomedicine* 2018;48:21–31. <https://doi.org/10.1016/j.phymed.2018.04.063>.
- [46] Arulseivan P, Tan W, Gothai S, Muniandy K, Fakurazi S, Esa N, et al. Anti-inflammatory potential of ethyl acetate fraction of *Moringa oleifera* in down-regulating the NF- κ B signaling pathway in lipopolysaccharide-stimulated macrophages. *Molecules* 2016;21:1452. <https://doi.org/10.3390/molecules-21111452>.
- [47] Fard M, Arulseivan P, Karthivashan G, Adam S, Fakurazi S. Bioactive extract from *Moringa oleifera* inhibits the pro-inflammatory mediators in lipopolysaccharide stimulated macrophages. *Phcog Mag* 2015;11:556. <https://doi.org/10.4103/0973-1296.172961>.
- [48] Wuryandari MRE, Widodo W, Widjajanto E, Rifa'i M. Activity red *Moringa oleifera* leaf extract as a preventive measure on the profile of CD4⁺CD62L⁺ and CD8⁺CD62L⁺ Cells in BALB/c mice injected *Salmonella typhimurium*. *KnE Soc Sci* 2019. <https://doi.org/10.18502/kss.v3i18.4727>.
- [49] Pérez-Cano F, Massot-Cladera M, Rodríguez-Lagunas M, Castell M. Flavonoids affect host-microbiota crosstalk through TLR modulation. *Antioxidants* 2014;3:649–70. <https://doi.org/10.3390/antiox3040649>.
- [50] Lee JK, Kim SY, Kim YS, Lee W-H, Hwang DH, Lee JY. Suppression of the TRIF-dependent signaling pathway of Toll-like receptors by luteolin. *Biochem Pharmacol* 2009;77:1391–400. <https://doi.org/10.1016/j.bcp.2009.01.009>.
- [51] Wuryandari MRE, Widodo N, Widjajanto E, Jatmiko YD, Rifa'i M. Red *Moringa oleifera* leaf fermentation extract protecting Hepatotoxicity in Balb/c mice injected with *Salmonella typhi* through Nrf-2, HO-1, and SOD-2 signaling pathways. *Res J Pharm Technol* 2020;13:1–6.
- [52] Chattopadhyay S, Sen GC. dsRNA-activation of TLR3 and RLR signaling: gene induction-dependent and independent effects. *J Interferon Cytokine Res* 2014;34:427–36. <https://doi.org/10.1089/jir.2014.0034>.
- [53] Sugiyama T, Kawaguchi K, Dobashi H, Miyake R, Kaneko M, Kumazawa Y. Quercetin but not luteolin suppresses the induction of lethal shock upon infection of mice with *Salmonella typhimurium*. *FEMS Immunol Med Microbiol* 2008;53:306–13. <https://doi.org/10.1111/j.1574-695X.2008.00398.x>.
- [54] Astry B, Venkatesha SH, Moudgil KD. Involvement of the IL-23/IL-17 axis and the Th17/Treg balance in the pathogenesis and control of autoimmune arthritis. *Cytokine* 2015;74:54–61. <https://doi.org/10.1016/j.cyt.2014.11.020>.
- [55] Wasanowati C, Sulistyansih E, Indradewa D, Kurniasih B. Morphophysiology and the yield of two types of moringa (*Moringa oleifera* Lamk) cultivated in two different regions in Madura. *IOP Conf Ser Earth Environ Sci* 2019;250: 12004. <https://doi.org/10.1088/1755-1315/250/1/012004>.
- [56] Noto Lana M, Sarnacki SH, Aya Castañeda M del R, Bernal MI, Giacomodonato MN, Cerquetti MC. Consumption of *Lactobacillus casei* fermented milk prevents *Salmonella* reactive arthritis by modulating IL-23/IL-17 expression. *PLoS One* 2013;8:e82588. <https://doi.org/10.1371/journal.pone.0082588>.
- [57] Matsumoto S, Hara T, Mitsuyama K, Nagaoka M, Tomiyasu N, et al. Probiotic *Lactobacillus*-induced improvement in murine chronic inflammatory bowel disease is associated with the down-regulation of pro-inflammatory cytokines in lamina propria mononuclear cells. *Clin Exp Immunol* 2005;140: 417–26. <https://doi.org/10.1111/j.1365-2249.2005.02790.x>.
- [58] Jaja-Chimedza A, Graf BL, Simmler C, Kim Y, Kuhn P, Pauli GF, et al. Biochemical characterization and anti-inflammatory properties of an isothiocyanate-enriched moringa (*Moringa oleifera*) seed extract. *PLoS One* 2017;12:e0182658. <https://doi.org/10.1371/journal.pone.0182658>.
- [59] Alqahtani WS, Albasher G. *Moringa oleifera* Lam. extract rescues lead-induced oxidative stress, inflammation, and apoptosis in the rat cerebral cortex. *J Food Biochem* 2021;45. <https://doi.org/10.1111/jfbc.13579>.
- [60] Belkaid Y, Tarbell K. Regulatory T cells in the control of host-microorganism interactions. *Annu Rev Immunol* 2009;27:551–89. <https://doi.org/10.1146/annurev.immunol.021908.132723>.
- [61] McArthur MA, Fresnay S, Magder LS, Darton TC, Jones C, Waddington CS, et al. Activation of *Salmonella typhi*-specific regulatory T cells in typhoid disease in a wild-type *S. Typhi* challenge model. *PLoS Pathog* 2015;11:e1004914. <https://doi.org/10.1371/journal.ppat.1004914>.
- [62] Goya-Jorge E, Jorge Rodriguez ME, Veitia MS-I, Giner RM. Plant occurring flavonoids as modulators of the aryl hydrocarbon receptor. *Molecules* 2021;26:2315. <https://doi.org/10.3390/molecules26082315>.
- [63] Mohammadi-Bardbori A, Bengtsson J, Rannug U, Rannug A, Wincent E. Quercetin, resveratrol, and curcumin are indirect activators of the aryl hydrocarbon receptor (AHR). *Chem Res Toxicol* 2012;25:1878–84. <https://doi.org/10.1021/tx300169e>.
- [64] Wang H-K, Yeh C-H, Iwamoto T, Satsu H, Shimizu M, Totsuka M. Dietary flavonoid naringenin induces regulatory T cells via an aryl hydrocarbon receptor mediated pathway. *J Agric Food Chem* 2012;60:2171–8. <https://doi.org/10.1021/jf204625y>.
- [65] Fallarino F, Gröhmman U, You S, McGrath BC, Cavener DR, Vacca C, et al. The combined effects of tryptophan starvation and tryptophan catabolites down-regulate T cell receptor ζ -chain and induce a regulatory phenotype in naive T cells. *J Immunol* 2006;176:6752–61. <https://doi.org/10.4049/jimmunol.176.11.6752>.
- [66] Xia Z-W, Zhong W-W, Xu L-Q, Sun J-L, Shen Q-X, Wang J-G, et al. Heme oxygenase-1-mediated CD4⁺ CD25⁺ high regulatory T cells suppress allergic airway inflammation. *J Immunol* 2006;177:5936–45. <https://doi.org/10.4049/jimmunol.177.9.5936>.
- [67] Atho'llah MF, Widyarti S, Rifa'i M. Elicited soybean (*Glycine max* L.) extract improves regulatory T cell activity in high fat-fructose diet mice. *Malang, Indonesia: AIP Publishing*; 2017. p. 20004. <https://doi.org/10.1063/1.4983415.1-020004-6>.

Lactobacillus plantarum FNCC 0137 fermented red Moringa oleifera exhibits protective effects in mice challenged with Salmonella typhi via TLR3/TLR4 inhibition and down-regulation of proinflammatory cy

ORIGINALITY REPORT

13%

SIMILARITY INDEX

15%

INTERNET SOURCES

10%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.researchgate.net Internet Source	7%
2	backend.orbit.dtu.dk Internet Source	2%
3	www.veterinaryworld.org Internet Source	2%
4	opac.wsb.torun.pl Internet Source	2%

Exclude quotes On

Exclude bibliography On

Exclude matches < 2%