

Identification of Pathogenic Bacteria in Liquid Waste from "X" Hospital in Kediri City

by Perpustakaan IIK Bhakti Wiyata

Submission date: 03-Mar-2025 11:07AM (UTC+0700)

Submission ID: 2501063034

File name: nal_of_Pharmaceutical_Sciences_2024_-_Binti_Mu_arofah_Kediri.pdf (179.11K)

Word count: 3273

Character count: 17863

Identification of Pathogenic Bacteria in Liquid Waste from "X" Hospital in Kediri City

Binti Mu'arofah¹, Tri Ana Mulyati^{2*}, Fery Eko Pujiono², Reza Alrayan²

¹Department of Medical Laboratory Technology, Institut Ilmu Kesehatan Bhakti Wiyata, Kediri, East Java, Indonesia

²Department of Pharmacy, Institut Ilmu Kesehatan Bhakti Wiyata, Kediri, East Java, Indonesia

Article history:

Submitted: 1-8-2024

Revised: 27-8-2024

Accepted: 5-9-2024

Corresponding author e-mail:
nanapujiono@gmail.com

Cite this article: Mu'arofah, B., Mulyati, T. A., Pujiono, F. E., Alrayan, R. (2024). Identification of Pathogenic Bacteria in Liquid Waste from "X" Hospital in Kediri City. Ad-Dawaa' J. Pharm. Sci. 7(2): 47-54.

Copyright:

This is an open-access article distributed under the terms of the CC BY-SA 4.0 license.



ABSTRACT

Introduction: Hospitals can not only be a place for people to seek treatment, but they can also get diseases, especially infectious diseases. One source of the spread of infectious diseases comes from hospital waste, so proper waste processing is needed. Hospital wastewater contains many pathogenic bacteria such as *Enterobacter cloacae*, *Bacillus* sp., *Kluyvera* sp., *Enterobacter sakazaki*, *Klebsiella oxytoca* and *E.coli*. This makes it necessary to manage hospital wastewater to prevent the spread of disease, especially by pathogenic bacteria. On the other hand, research is still needed on the specific types of bacteria found in wastewater, including hospitals in Kediri City. **Aims:** to determine the types of pathogenic bacteria at the "X" hospital in Kediri City. **Result:** the density of bacteria in the bottom waste was greater than in the top and middle parts, with an average of 1,830 bacteria/100 ml. The results of the identification test showed that pathogenic bacteria contained in the waste from hospital "X" in Kediri City included *Klebsiella* (25%), *E.coli* (50%) and *Pseudomonas* (25%). **Conclusion:** The waste from "X" Hospital Kediri City contains three types of pathogenic bacteria: *Klebsiella*, *E. coli*, and *Pseudomonas*.

KEYWORDS: Bacteria, waste, hospital, Kediri

INTRODUCTION

A hospital is a health installation to care for and treat various diseases. On the other hand, hospitals are also places where people get different kinds of diseases, especially infectious diseases. One source of the spread of infectious diseases comes from hospital waste. Waste is the remainder of the activities of living creatures originating from various activities and can be in solid, liquid, or gas form (Singh, 2023). Hospital waste is all the waste generated from hospital activities that

can pollute the environment and potentially transmit infections (Ningrum et al., 2017). Generally, hospital waste is divided into medical and non-medical waste. Based on its form, hospital waste can be categorized into three types: solid waste, liquid waste, and gas waste (Chand et al., 2020). One type of dangerous hospital waste is hospital liquid waste.

Hospital liquid waste is the remaining waste product or all wastewater, including feces originating from hospital activities, which

B. Mu'arofah, et al.

contains ¹⁰ pathogenic bacteria such as *Escherichia coli*, *Bacillus subtilis*, *Staphylococcus aureus*, and *Klebsiella pneumoniae*. Hospital liquid waste that is not managed correctly can cause the spread of infectious diseases. Therefore, hospitals must manage liquid waste according to applicable requirements and standards (Bhandari et al., 2023). Liquid waste quality is the condition of liquid waste, expressed in terms of discharge, content, and pollutant materials. The maximum discharge by Ministry of Environment Regulation Number Kep/51/MENLH/10/1995 is the highest discharge that is still permitted to be discharged into the environment. Important liquid waste quality parameters to know are ⁷ suspended solids, dissolved solids, Biochemical Oxygen Demand (BOD), Chemical Oxygen Demand (COD), coliform organisms, pH, Dissolved Oxygen (DO), chlorine requirements, nutrients, ²⁵ heavy metals, and other parameters (Ranti et al., 2020).

The increased number of healthcare facilities in hospitals causes increased environmental pollution. This also causes an increased risk of spreading infectious diseases. The research results of Astawa & Tarini (2017) show that wastewater at Sanglah Hospital contains several pathogenic bacteria such as *Enterobacter cloacae*, *Bacillus sp.*, *Enterobacter Okazaki*, *Klebsiella oxytoca*, and *E.coli*. Similar results were also reported by Winarti (2020), who reported that the wastewater at RSKIA Sadewa contained Total

Coliform bacteria of 2400×10^3 MPN/100ml. This shows that it is necessary to manage hospital wastewater to prevent the spread of diseases, especially those caused by pathogenic bacteria.

¹² Based on the results of this research, it is necessary to identify bacteria in hospital waste, including hospital "X" in Kediri City. This study aims to identify the types of bacteria still present in the waste from hospital "X" in Kediri City. This research is essential to prevent the transmission of infectious diseases due to hospital waste contamination, control nosocomial infections and antibiotic resistance, support better health policies, and improve preparedness and response to potential disease outbreaks.

METHODS

Materials

The materials needed in this study are Lakosa Broth 1 (LB I), Lakosa Broth 3 (LB III), Brilian ²⁰ Green Lactose Broth (BGLB), Eosin Methylene Blue (EMB), Centrimid, Mannitol Slat agar (MSA), Salmonella Shigella Agar (SSA), Tripel Suger Iron Agar (TSIA) and Indol, Methil red, Voges Prouskover, Citrat (IMVC).

Preparation and Sampling

Sterile glass bottles are opened at 120°C for 1 hour. Waste collection is carried out at the inlet and outlet of the IPAL (outlet) at the "X" hospital in Kediri City, with the number of samples in this study being 12 samples (2

IPALs, each IPAL is taken at 3 points, namely the top, middle, bottom points and repeated two times). The samples were 100ml of outlet liquid waste per bottle; samples were taken within a day using sterile bottles coded and taken to the laboratory with a dry ice box for examination, with standardized parameters (Yatini, et. al., 2021).

Bacterial Identification

Preliminary test

A total of 5 tubes containing LB III were added, each with a sample of hospital waste "X" 10 ml. Five LB I tubes were prepared and filled with 1ml of sample each. Each tube was then taken in 0.1 ml (100 ul), and all treatments were carried out sterily in the LAF. The tubes filled with water samples were homogenized so that the waste samples were evenly mixed with the LB III and LB I media; LB III and LB I media were incubated in an incubator at 37 °C for 24 hours (Yatini, et. al., 2021).

Confirmation test

Observing changes in each LB III and LB I tube that had been incubated to see the turbidity and the emergence of gas. If Media LB III and I are vivacious, the confirmation test is continued by inoculating the sample into the BGLB media in a sterile manner in LAF. The results are incubated in an incubator for 24 hours at a temperature of 37°C (Yatini, et. al., 2021).

Complementary test

We are observing changes in each BGLB tube incubated for 24 hours. Positive results

Identification of Pathogenic bacteria are indicated by samples that turn cloudy and form gas. The results in the confirmation test tube are adjusted to the MPN (Most Probable Number) table of variety 555. Positive BGLB results are then inoculated into EMB, Centrimtde, SSA, MSA, TSIA, and IMVC media and then incubated for 24 hours at a temperature of 37°C to identify bacteria in the waste of the hospital "X" Kediri City (Yatini, et. al., 2021).

RESULTS AND DISCUSSION

This study identified pathogenic bacteria in the waste of hospital "X" Kediri City. The study identified bacteria at 3 points: the top, middle, and bottom. The results of bacterial density at various sampling points are shown in Table. 1. The results of bacterial identification in the waste of hospital "X" Kediri City are shown in Table 2.

Table 1 shows that the waste of hospital "X" Kediri City at the bottom has more bacteria than at the top and bottom. The density of the bacteria in the lower waste shows an average of 1,830 bacteria/100 ml. Several things can affect the higher density of bacteria in the upper part: environmental conditions, nutrients, and bacterial sedimentation (Fletcher, 2015). At the bottom of the waste reservoir, it generally has a lower temperature and high humidity so that bacteria can grow more easily. In addition, at the bottom of the waste container, there is a possibility of sedimentation of organic materials from hospital waste (such as blood and tissue) so

Table 1. Bacterial Density at Various Sampling Points in The Waste of Hospital "X" Kediri City

No	Sample code	Positive Combination	Bacterial Density/100 ml
1	A1.1	5-5-2	540
2	A1.2	5-5-2	540
3	A2.1	5-5-5	>2400
4	A2.2	5-5-4	1600
5	A3.1	5-5-5	>2400
6	A3.2	5-5-5	>2400
7	B1.1	5-4-2	220
8	B1.2	5-4-4	350
9	B2.1	5-5-2	540
10	B2.2	5-5-2	540
11	B3.1	5-5-4	1600
12	B3.2	5-5-3	920

A shelter, 8 top dot, 2 middle dots, 3 bottom dots, repeat 2x
 B shelter, 1 top dot, 2 middle dots, 3 bottom dots, repeat 2x

that it becomes a source of nutrition for bacterial growth (Mahdi & Gomes, 2019). The density of bacteria also tends to be higher because piles of solid particles generally settle at the bottom of the waste (Maina & Susan Muthoni, 2020).

Table 2 shows where the waste from the hospital "X" in Kediri City contains 3 types of pathogenic bacteria, namely *Klebsiella* bacteria 25%, *E. coli* 50%, and *Pseudomonas* 25%. These results are by the research of Cuetero-Martínez et al. (2023) which identified the presence of pathogens in hospital waste: *Escherichia coli*, *Enterobacter spp.*, *Pseudomonas aeruginosa*, and *Bacillus spp.* Zhang et al. (2021) also reported similar results and identified various pathogenic bacteria, including *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus* in hospital waste. The presence of bacteria in hospital waste is due to the physiological properties of pathogenic bacteria and resistance to disinfectants and

antibiotics, contributing to their presence in hospital waste Gönder et al. (2021).

Table 2 shows that the type of pathogenic bacteria most often found in hospital waste "X" in Kediri City is *E. coli*. *E. coli* is the main cause of hospital urinary tract infections (UTIs) and gastrointestinal infections. Many hospital patients are treated for this condition, so hospital wastewater generally contains pathogenic *E. coli* bacteria from patients with this infection (Osińska et al., 2017). In addition, *E. coli* can survive in various environmental conditions, including in-hospital wastewater, which often has varying pH and changing nutrient concentrations, and can also utilize various organic substrates in hospital waste, including nutrients from food waste and metabolic products of the body (Akther et al., 2018; Paulshus et al., 2019). These results cause *E. coli* in hospital wastewater, including Hospital "X" in Kediri City.

Identification of Pathogenic bacteria

Table 2. Bacterial Identification in The Waste of Hospital "X" Kediri City

Sample code	Colour of Colony Selective Media	TSIA	Indol	MR	VP	Citrat	Bacterial identification
A1.1	<i>Metalic sheen</i>	L= acid, D=acid, H2S = -, Gas = +	+	+	-	-	<i>E. coli</i>
A1.2	<i>Pink Mucoïd</i>	L= acid, D=acid, H2S = -, Gas = +	-	-	+	+	<i>Klebsiella</i>
A2.1	<i>Metalic sheen</i>	L= acid, D=acid, H2S = -, Gas = +	+	+	-	-	<i>E. coli</i>
A2.2	<i>Pink Mucoïd</i>	L= acid, D=acid, H2S = -, Gas = +	-	-	+	+	<i>Klebsiella</i>
A3.1	<i>Metalic sheen</i>	L= acid, D=acid, H2S = -, Gas = +	+	+	-	-	<i>E. coli</i>
A3.2	<i>Pink Mucoïd</i>	L= acid, D=acid, H2S = -, Gas = +	-	-	+	+	<i>Klebsiella</i>
B1.1	<i>Metalic sheen</i>	L= acid, D=acid, H2S = -, Gas = +	+	+	-	-	<i>E. coli</i>
B1.2	<i>Phyocianin</i>	L= alkalis, D=alkalis, H2S = -, Gas = -	-	-	-	+	<i>Pseudomonas</i>
B2.1	<i>Metalic sheen</i>	L= acid, D=acid, H2S = -, Gas = +	+	+	-	-	<i>E. coli</i>
B2.2	<i>Phyocianin</i>	L= alkalis, D=alkalis, H2S = -, Gas = -	-	-	-	+	<i>Pseudomonas</i>
B3.1	<i>Metalic sheen</i>	L= acid, D=acid, H2S = -, Gas = +	+	+	-	-	<i>E. coli</i>
B3.2	<i>Phyocianin</i>	L= alkalis, D=alkalis, H2S = -, Gas = -	-	-	-	+	<i>Pseudomonas</i>

Selective media used:

Eosin Methylene Blue (EMB) for *E.coli* and *Klebsiella* bacteria.

Centrimed for *Pseudomonas* bacteria

Table 2 also shows that besides *E. coli*, two other pathogenic bacteria are found in Hospital "X" in Kediri City, namely *Klebsiella* and *Pseudomonas*. *Klebsiella spp.* is a gram-negative bacteria known as an opportunistic pathogen. *Klebsiella spp.* has a polysaccharide capsule that helps evade the immune system

and increases its virulence. This capsule also helps bacteria survive in hostile environments, such as hospital waste (Gomes et al., 2021). *Klebsiella spp.* can also utilize various carbon sources, including glucose and lactose, which are common in hospital waste and survive in low oxygen conditions. This causes *Klebsiella*

B. Mu'rofah, et al.

spp. to be often found in hospital waste, including in the waste of hospital "X" in Kediri City (Sakkas et al., 2019).

Another type of pathogenic bacteria found in the waste of hospital "X" in Kediri City is *Pseudomonas*. *Pseudomonas aeruginosa* is a highly adaptive bacteria and can survive in various environmental conditions, including environments with high humidity and inconsistent nutrients, such as hospital waste (Khan et al., 2024). One of the main reasons why *Pseudomonas aeruginosa* is often found in hospital waste is its ability to form biofilms (Divyashree et al., 2022). Biofilms are communities of bacteria that adhere to surfaces and are protected by an extracellular matrix, which makes them more resistant to cleaning and disinfection. Hospital waste often has a variety of surfaces that can support biofilm formation (Suleyman et al., 2018).

CONCLUSION

Based on the research results, it can be concluded that the density of the number of bacteria in the lower part of the waste is greater than in the upper and middle parts, with an average of 1,830 bacteria / 100 ml. For identification results, *Klebsiella* bacteria were obtained from 3 samples (25%), *E. coli* from 6 samples (50%), and *Pseudomonas* from 3 samples (25%).

ACKNOWLEDGEMENT

The author would like to thank the Ministry of Education, DIRJEN DIKTI, for providing funding through the "Penelitian Dosen

Pemula" grant. The author would also like to thank the Bhakti Wiyata Foundation and the Bhakti Wiyata Institute of Health Sciences for supporting the implementation of this research. The author would also like to thank the D3 Medical Laboratory Technology and S1 Pharmacy students who have helped the research process.

REFERENCES

- Akther, S., Debnath, T., & Chowdhury, M. M. H. (2018). Multidrug resistant *E. coli* in hospital waste water: a potential concern for public health. *Adv. Biotechnol. Microbiol.*, 8(55572910.19080). <https://doi.org/10.19080/aibm.2018.08.55729>
- Astawa, I. B. B., & Tarini, N. M. A. (2017). Identifikasi Jenis Bakteri dalam Air Limbah di Rumah Sakit Sanglah. *E-Jurnal Medika*, 6(6), 1–6.
- Bhandari, G., Chaudhary, P., Gangola, S., Gupta, S., Gupta, A., Rafatullah, M., & Chen, S. (2023). A review on hospital wastewater treatment technologies: Current management practices and future prospects. *Journal of Water Process Engineering*, 56, 104516. <https://doi.org/10.1016/j.jwpe.2023.104516>
- Chand, S., Shastry, C. S., Hiremath, S., Joel, J. J., Bhat, C. K., & Mateti, U. V. (2020). Water, sanitation, hygiene and biomedical waste disposal in the healthcare system: A review. *Biomedicine*, 40(1), 14-19.
- Cuetero-Martínez, Y., de Los Cobos-Vasconcelos, D., Aguirre-Garrido, J. F., Lopez-Vidal, Y., & Noyola, A. (2023). Next-generation sequencing for surveillance of antimicrobial resistance and pathogenicity in municipal wastewater treatment plants. *Current Medicinal Chemistry*, 30(1), 5–29. <https://doi.org/10.2174/0929867329666220802093415>
- Divyashree, M., Mani, M. K., & Karunasagar, I. (2022). Association of exopolysaccharide genes in biofilm deve-

- loping antibiotic-resistant *Pseudomonas aeruginosa* from hospital wastewater. *Journal of Water and Health*, 20(1), 176–184. <https://doi.org/10.2166/wh.2021.223>
- Fletcher, S. (2015). Understanding the contribution of environmental factors in the spread of antimicrobial resistance. *Environmental Health and Preventive Medicine*, 20(4), 243–252. <https://doi.org/10.1007/s12199-015-0468-0>
- Gomes, A. É. I., Pacheco, T., Santos, C. da S. dos, Pereira, J. A., Ribeiro, M. L., Darrieux, M., & Ferraz, L. F. C. (2021). Functional insights from KpfR, a new transcriptional regulator of fimbrial expression that is crucial for *Klebsiella pneumoniae* pathogenicity. *Frontiers in Microbiology*, 11, 601921. <https://doi.org/10.3389/fmicb.2020.601921>
- Gönder, Z. B., Kara, E. M., Celik, B. O., Vergili, I., Kaya, Y., Altinkum, S. M., Bagdatli, Y., & Yilmaz, G. (2021). Detailed characterization, antibiotic resistance and seasonal variation of hospital wastewater. *Environmental Science and Pollution Research*, 28, 16380–16393. <https://doi.org/10.1007/s11356-020-12221-w>
- Khan, M. A. S., Islam, Z., Shah, S. M. T., & Rahman, S. R. (2024). Characterization of biofilm formation and multi-drug resistance among *Pseudomonas aeruginosa* isolated from hospital wastewater in Dhaka, Bangladesh. *Journal of Water and Health*, 22(5), 825–834. <https://doi.org/10.2166/wh.2024.294>
- Mahdi, A. B., & Gomes, C. (2019). Effects of microwave radiation on micro-organisms in selected materials from healthcare waste. *International Journal of Environmental Science and Technology*, 16(3), 1277–1288. <https://doi.org/10.1007/s13762-018-1741-8>
- Maina, & Susan Muthoni. (2020). *Isolation of Bacteria and Prevalence of Resistant Genes in Isolates from Hospital Waste and Surfaces and its Management in Hospitals in Kenya*. Jomo Kenyatta University of Agriculture and Technology.
- Ningrum, P. T., & Khalista, N. N. (2017). Gambaran Pengelolaan Limbah Cair Di Rumah Sakit X Kabupaten Jember. *Jurnal Ilmu Kesehatan Masyarakat*, 10(2).
- Osińska, A., Korzeniewska, E., Harnisz, M., & Niestępski, S. (2017). The prevalence and characterization of antibiotic-resistant and virulent *Escherichia coli* strains in the municipal wastewater system and their environmental fate. *Science of the Total Environment*, 577, 367–375. <https://doi.org/10.1016/j.scitotenv.2016.10.203>
- Paulshus, E., Kühn, I., Möllby, R., Colque, P., O'Sullivan, K., Midtvedt, T., Lingaas, E., Holmstad, R., & Sørum, H. (2019). Diversity and antibiotic resistance among *Escherichia coli* populations in hospital and community wastewater compared to wastewater at the receiving urban treatment plant. *Water Research*, 161, 232–241. <https://doi.org/10.1016/j.watres.2019.05.102>
- Ranti, J. C. A., Sutanto, H. B., & Prihatmo, G. (2020). Efektivitas Sistem Constructed Wetland Dalam Menurunkan Jumlah Bakteri Patogen Pada Limbah Rumah Sakit. *SAINTEK: Jurnal Ilmiah Sains Dan Teknologi Industri*, 4(2), 89–93. <https://doi.org/10.32524/sainstek.v4i2.160>
- Sakkas, H., Bozidis, P., Ilia, A., Mpekoulis, G., & Papadopoulou, C. (2019). Antimicrobial resistance in bacterial pathogens and detection of carbapenemases in *Klebsiella pneumoniae* isolates from hospital wastewater. *Antibiotics*, 8(3), 85. <https://doi.org/10.3390/antibiotics8030085>
- Singh, D. (2023). Advances in industrial waste management. In *Waste management and resource recycling in the developing world* (pp. 385–416). Elsevier. <https://doi.org/10.1016/B978-0-323-90463-6.00027-0>
- Suleyman, G., Alangaden, G., & Bardossy, A. C. (2018). The role of environmental contamination in the transmission of nosocomial pathogens and healthcare-

B. Mu'rofah, et al.

associated infections. *Current Infectious Disease Reports*, 20, 1–11.

Winarti, C. (2020). Penurunan bakteri total coliform pada air limbah rumah sakit terhadap pengaruh lama waktu penyinaran dengan sinar ultra violet. *Jurnal Rekayasa Lingkungan*, 20(1).
<https://doi.org/10.1007/s11908-018-0620-2>

Yatini, Y., & Harahap, A. (2021). The Density of Coliform Bacteria Indicators of Pollution in the River Right Labuhanbatu Selatan. *Budapest International Research*

and Critics Institute (BIRCI-Journal) Humanities and Social Sciences, 4(1), 1248-1253.

<https://doi.org/10.33258/BIRCI.V4I1.1744>

Zhang, S., Li, X., Wu, J., Coin, L., O'brien, J., Hai, F., & Jiang, G. (2021). Molecular methods for pathogenic bacteria detection and recent advances in wastewater analysis. *Water*, 13(24), 3551.

<https://doi.org/10.3390/w13243551>

Identification of Pathogenic Bacteria in Liquid Waste from "X" Hospital in Kediri City

ORIGINALITY REPORT

13%	10%	9%	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	www.researchgate.net Internet Source	2%
2	jnk.phb.ac.id Internet Source	1%
3	wjarr.com Internet Source	1%
4	Dody Karnovel, John Hendri, Alfian Asri. "Physiological Response of Broiler Chicken To Cage Temperature", Journal of Animal Nutrition and Production Science, 2023 Publication	1%
5	idus.us.es Internet Source	1%
6	journals.plos.org Internet Source	1%
7	nepis.epa.gov Internet Source	1%
8	Glass, R.I.. "Nucleotide sequence of the structural glycoprotein VP7 gene of Nebraska calf diarrhea virus rotavirus: Comparison with homologous genes from four strains of human and animal rotaviruses", Virology, 198503 Publication	1%
9	Mehdi Zarei, Mohsen Paknejad, Mohammad Hadi Eskandari. "Sublethal chlorine stress	1%

promotes the biofilm-forming ability of Salmonella enterica serovars enteritidis and expression of the related genes", Food Microbiology, 2023

Publication

10

www.bibliomed.org

Internet Source

<1 %

11

iris.who.int

Internet Source

<1 %

12

www.ncbi.nlm.nih.gov

Internet Source

<1 %

13

Ely Isnaeni, Erna Rahmawati. "Moringa Nastar Increase Prolaktin As Support Of 1000 HPK Action With UNICEF Consept Approach", Journal Of Nursing Practice, 2019

Publication

<1 %

14

Gisele Canan-Rochenbach, Marco A.B. Barreiros, André O.S. Lima, Aline Dal Conti-Lampert et al. "Are hospital wastewater treatment plants a source of new resistant bacterial strains?", Environmental Science and Pollution Research, 2023

Publication

<1 %

15

Markus Kaban, Tri Retnaningsih Soeprowati, Suhartono Suhartono, Purwanto Adi Pireno et al. "Informing compliance factors regarding hospitals waste management procedures in Indonesia: a scoping review", International Journal of Public Health Science (IJPHS), 2025

Publication

<1 %

16

Sardar Karash, Young Min Kwon. "Iron-dependent essential genes in Salmonella

<1 %

Typhimurium", Cold Spring Harbor
Laboratory, 2018

Publication

17 W. J. Barden, A. W. Sargent. "The Lake
Washington Ship Canal, Washington",
Transactions of the American Society of Civil
Engineers, 1928 <1 %
Publication

18 journalseeker.researchbib.com <1 %
Internet Source

19 www.earthdoc.org <1 %
Internet Source

20 www.scribd.com <1 %
Internet Source

21 Ana Érika Inácio Gomes, Thaisy Pacheco,
Cristiane da Silva dos Santos, José Aires
Pereira et al. "Functional Insights From KpfR, a
New Transcriptional Regulator of Fimbrial
Expression That Is Crucial for Klebsiella
pneumoniae Pathogenicity", Frontiers in
Microbiology, 2021 <1 %
Publication

22 perpustakaan.poltekkes-malang.ac.id <1 %
Internet Source

23 studentsrepo.um.edu.my <1 %
Internet Source

24 www.neliti.com <1 %
Internet Source

25 "Green Technologies for Industrial Waste
Remediation", Springer Science and Business
Media LLC, 2023 <1 %
Publication

26

Maria Eduarda Souza Guerra, Giulia Destro, Brenda Vieira, Alice S. Lima et al. "Klebsiella pneumoniae Biofilms and Their Role in Disease Pathogenesis", *Frontiers in Cellular and Infection Microbiology*, 2022

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On